

## First report of *Alternaria alternata* associated with seed of Asafetida (*Ferula assa-foetida*)

Nima Khaledi<sup>1\*</sup>, Farshid Hassani<sup>1</sup>

### Abstract

*Alternaria* is considered one of the most common saprophytic fungi as well as is recognized as pathogens of various plant parts including seed. In May 2020, chlorosis and blight symptoms observed on seedlings emerging from seeds of Asafetida (*Ferula assa-foetida*). Using the agar plate test, a total of 19 fungal isolates were isolated from seed samples collected from different Asafetida growing regions of Iran. Based on morphological and molecular characteristics, all the isolates was identified as *Alternaria alternata*. Pathogenicity tests were conducted and Koch's postulates were fulfilled by re-isolation of the identified fungus. This is the first report on isolation and identification of *A. alternata* associated with seed of Asafetida in world.

**Key words:** *Alternaria alternata*, Asafetida, Germination test, Pathogenicity, Seed-borne.

---

<sup>1</sup> Seed and Plant Certification and Registration Research Institute, Agricultural Research, Education and Extension Organization (AREEO), Karaj, Iran. \* Corresponding author: Email: n\_khaledi@areeo.ac.ir

## Introduction

Asafetida (*Ferula assa-foetida* belonging Apiaceae family), is an important spice crop with medicinal properties and native to central Asia, eastern Iran to Afghanistan. In previous studies, antioxidant (Kavoosi and Rowshan 2013; Dehpour et al., 2009), antimicrobial (Bhatnager et al., 2015; Divya et al., 2014), antifungal (Rani et al., 2009; Kamble and Patil, 2008) and antiviral (Mahendra and Bisht 2012) effects of this plant have been demonstrated. To date, no fungal pathogens have been reported on *F. assa-foetida*.

## Materials and methods

In May 2020, samples of 400 seeds from each of Asafetida seed crops in Shiraz, Mashhad, Isfahan and Yasuj, Iran were tested for germination on moistened paper in petri dishes. The germination test was conducted in four replications of 100 seeds each sample by using paper (between papers) medium and was incubated at  $25 \pm 1$  °C temperature and  $85 \pm 5$  % relative humidity for 14 d.

Different seed health testing methods viz standard blotter and agar plate tests were used to investigate the fungal pathogens

present in these seeds and their effect on germination (ISTA 2013). The seeds were surface-sterilized in 1% sodium hypochlorite solution for one minute and rinsed with sterile distilled water. The collected seeds samples were plated in petri dishes containing sterile wet filter paper or potato dextrose agar (PDA) medium and incubated for 7 days at  $25 \pm 1$  °C under alternating cycles of 12 hours light and darkness. Among different Asafetida samples in the investigated regions, fungi-infected seeds was detected only in sample Shiraz.

The emerged fungi were picked up, purified and identified based on morphological and molecular characteristics. Pure cultures were obtained by using single spore and hyphal tip methods at  $23 \pm 1$  °C on 2% water agar (2% WA) and 0.05% potato dextrose agar (0.05% PDA) media. For morphological identification, the isolates were cultured on Potato Carrot Agar medium (PCA) and incubated at  $25 \pm 1$  °C under cool-white fluorescent illumination with 16 h dark, 8 h light photoperiod for 14 d (Simmons, 2007). Preliminary identification of fungal isolates was carried out by using standard keys as described by Barnett and Hunter (1998), and Simmons (2007).

Further confirmation of the identification was obtained by molecular characterization in which genomic DNA was extracted using DNA extraction kit (Genomic DNA isolation kit IV; DENA Zist Asia, Iran) according to the manufacturer's instructions. To verify the identity, the internal transcribed spacer (ITS) of rDNA was amplified and sequenced using primers ITS1 (5'-TCCGTAGGTGAACCTGCGG-3') and ITS4 (5'-TCCTCCGCTTATTGATATGC-3'), specifically designed for *A. alternata* (Zheng et al., 2015) and sequenced by Macrogen (Korea). Polymerase chain reaction (PCR) products were sequenced and the identity of fungal isolates confirmed by a BLAST search on the GenBank database. PCR identification for the isolate was confirmed using AaltFor (5'-GTGCCTTCCCCAAGGTCTCCG-3') and AaltRev (5'-CGGAAACGAGGTGGTTCAGGTC-3') primer specific as the *A. alternata* which produced fragment size of approximately 184 base pair (bp) (Kordalewska et al., 2015).

In order to determinate if *A. alternata* was the possible cause of discolored seeds and abnormal seedlings and also to confirm Koch's postulates, the fungal isolate was subjected to pathogenicity tests on seeds and

seedlings. Asafetida seeds were inoculated with a conidial suspension ( $1 \times 10^5$  conidia  $\text{mL}^{-1}$  amended with 0.05% Tween 20) and incubated for 10 d at  $25 \pm 2$  °C in a growth chamber under 12 h photoperiod. For pathogenicity test on seedlings, 1000  $\mu\text{l}$  of spore suspension ( $1 \times 10^5$  conidia  $\text{mL}^{-1}$  amended with 0.05% Tween 20) was sprayed onto the leaves of seedlings at the two- to three leaf stage. Plants were covered by polyethylene bags for 48 hours, maintained in a greenhouse at  $25 \pm 2$  °C and 90-100% humidity. After 72 hours of incubation, the appeared chlorosis and blighting developed on the leaves of inoculated plants, whereas control plants remained asymptomatic. The pathogenicity tests were repeated twice under the same conditions. After germinating the seeds, 19% of the resulting seedlings were abnormal (deformed and diseased), discolored, and most of them (average of 17% of all seedlings) died shortly after emergence. The mean frequency of fungi recorded was found to be significantly highest in agar plate test (19%) when compared to standard blotter test (15%). All the 19 fungal isolates isolated were similar in morphological features.

## Results and Discussion

The fungi isolated were initially light grey in color and changed to dark green, and then the whole upper surface of the colony turned black after 3 days of incubation on PDA medium. The color of the colony was gray-olive green on PCA medium. Conidiophores that developed singly or in clusters, were light brown to brown, were long or short being either ramified or simple and measured 33-94  $\mu\text{m}$  in length and 4-10

$\mu\text{m}$  in width. Conidia were produced in acropetal chains on conidiophores, light olivaceous to dark brown, varied in shape from ovoid to ellipsoidal with 4-7 transverse septa and 0-3 longitudinal septa, and measured 18-30  $\mu\text{m}$  in length and 6-11  $\mu\text{m}$  in width (Fig. 1). Based on morphological characterization, the fungi isolated from seed were tentatively identified as *A. alternata* (Simmons, 2007).




Fig. 1. *Alternaria alternata* isolate AANKH 568 from 10-day-old colony on PCA (a) conidiophores with conidia developing at the tips, (b, c) mature conidia, (d) germinating conidium. Scale bars 20  $\mu\text{m}$ .

Ten days after inoculation, the seedlings inoculated with fungal isolate showed chlorosis and blight symptoms. The symptoms on the tissues resembled those observed in the field. No symptoms were detected on the controls. *A. alternata* was recovered from the diseased tissues but not from the control. Fungal isolate recovered from seedling was identified based on morphological characteristics including cultural morphology, size and shape of conidia, and sporulation patterns and compared with the initial isolate. This combination of data confirmed that the pathogen was *A. alternata*.

*A. alternata* was re-isolated from the seeds and seedling exhibited the similar morphology and molecular of the earlier described one and also proving pathogenicity and confirming Koch's postulates. The results showed that the *A. alternata* causes seed discoloration, shrivelling, reduction in seed germination and increase in the number of abnormal (deformed and diseased) seedlings. The isolate of *A. alternata* has been lodged in the culture collection of the Seed Health Laboratory, the Seed and Plant Certification

and Registration Institute (SPCRI) in Iran with the accession number AANKH 568.

The fungus *A. alternata* is one of the most common pathogens and usually reduce the quality of seed that experience stress during the ripening period, particularly from excessive rain and delayed harvest. This *Alternaria*-infected seeds is may be attributed to cultures of the sensitive native population, varying climate and environmental conditions, crop rotation with vegetables and also sensitive crops in the Shiraz region of Iran. Further studies should be considered to determine potential yield loss caused by *A. alternata*, as well as control strategies to limit the spread of this pathogen. However, some management strategies, such as crop rotation with non-host crops, field sanitation, seed treatment with fungicides and use of healthy seeds may be helpful reducing disease severity. *A. alternata* is seed-borne and seed-transmitted pathogen in Asafetida, and it is introduced into new regions through the movement of *A. alternata*-infected seed. To our knowledge, this is the first report of *A. alternata* has been isolated from Asafetida seeds produced in world.

## References

- [ISTA] International Seed Testing Association (2013) The germination test. In: "International Rules for Seed Testing". ISTA, Bassersdorf, Switzerland
- Barnett HL, Hunter BB, (1998) "Illustrated Genera of Imperfect Fungi" 4th ed. Minneapolis, MN, Burgess
- Bhatnager R, Rani R, Dang AS, (2015) "Antibacterial activity of *Ferula asafoetida*: a comparison of red and white type" J Appl Biol Biotechnol 3:18-21.
- Dehpour AA, Ebrahimzadeh MA, Fazel NS, Mohammad NS, (2009) "Antioxidant activity of the methanol extract of *Ferula asafoetida* and its essential oil composition" Grasas Aceites 60:405-412.
- Divya K, Ramalakshmi K, Murthy PS, Rao LJM, (2014) "Volatile oils from *Ferula asafoetida* varieties and their antimicrobial activity" LWT Food Sci Technol 59:774-779.
- Kamble VA, Patil SD, (2008) "Spice-derived essential oils: effective antifungal and possible therapeutic agents" J Herbs Spices Med Plants 14:129-143.
- Kavoosi G, Rowshan V, (2013) "Chemical composition, antioxidant and antimicrobial activities of essential oil obtained from *Ferula asafoetida* oleo-gum-resin: effect of collection time" Food Chem 138:2180-2187.
- Kordalewska M, Brillowska-Dąbrowska A, Jagielski T, Dworecka-Kaszak B, (2015) "PCR and real-time PCR assays to detect fungi of *Alternaria alternata* species" Acta Biochim Pol 62:707-712.
- Mahendra P, Bisht S, (2012) "*Ferula asafoetida*: traditional uses and pharmacological activity" Pharmacogn Rev 6:141-146.
- Rani A, Jain S, Dureja P, (2009) "Synergistic fungicidal efficacy of formulations of neem oil, nicotinic acid and *Ferula asafoetida* with  $\alpha$ ,  $\beta$ -unsaturated carbonyl compounds against *Sclerotium rolfsii* ITCC 5226 & *Macrophomina phaseolina* ITCC 0482" J Pestic Sci 34: 253-258.
- Simmons EG, (2007) *Alternaria*. An identification manual. In: CBS biodiversity series 6CBS Fungal Biodiversity Centre, Utrecht, The Netherlands
- Zheng HH, Zhao J, Wang TY, Wu XH, (2015) "Characterization of *Alternaria* species associated with potato foliar diseases in China" Plant Pathol 64:425-433.